

Introduction

- Discovered in the slope eastern coast of New Caledonia in 1996 by Pietra and coworkers from a new genus of calcareous sponge, *Leudcasandra caveolata*.
- Highly potent cytotoxic against KB oral epidermoid carcinoma ($IC_{50} = 0.05 \mu\text{g mL}^{-1}$) and P388 leukemia ($IC_{50} = 0.25 \mu\text{g mL}^{-1}$) cell lines as well as a powerful inhibitor of *Candida albicans*, a pathological yeast that attacks AIDS patient.
- Biogenesis origin unknown.
- Suspected to be of microbial origin.
- Later expedition failed to collect any more material.

Helv. Chim. Acta **1996**, 79, 51.

Helv. Chim. Acta **1999**, 82, 347.


Introduction

■ Structural features



- Doubly O-bridged 18-membered macrolactone ring structure
- Extensive 1,3-dioxygenation
- Unusual oxazole bearing side chain

■ Synthetic interest stirred by its unavailability, high bioactivity and unique structure


■ Up-to-date, there are 2 formal syntheses and 4 total syntheses reported



Retrosynthetic Analyses





Leighton, J. L. *et al. J. Am. Chem. Soc.* **2000**, *122*, 12894.





Rychnovsky, S. D. *et al. J. Am. Chem. Soc.* **2001**, *123*, 8420.

Retrosynthetic Analyses




Wipf, P. *et al. Chem. Commun.* **2002**, 2006.
Wipf, P. *et al. J. Org. Chem.* **2001**, 66, 3242.




Kozmin, S. A. *et al. J. Am. Chem. Soc.* **2002**, 124, 13670.
Kozmin, S. A. *Org. Lett.* **2001**, 3, 755.

Retrosynthetic Analyses




Carreira, E. M. *Angew. Chem. Int. Ed.* **2002**, *41*, 2908.





Paterson, I. *et al. Angew. Chem. Int. Ed.* **2003**, *42*, 343.


Carreira Synthesis: Construction of C1-C9 Fragment



42 %
91 % ee




Carreira Synthesis: Construction of C1-C10 Fragment




63 %
dr = 10:1

TBSCl, imidazole,
DMAP, DMF, rt





96 %

20 mol %
PdCl₂,
CuCl, O₂,
DMF/H₂O = 7:1,
rt





86 %


Carreira Synthesis: Construction of C11–C18 fragment




Carreira Synthesis: Coupling of Two Main Fragments






Carreira Synthesis: Macrolactonization






Carreira Synthesis: Installation of Side Chain C19-C22



Carreira/Leighton Synthesis: Side Chain C2'-C11'



Carreira/Leighton: Completion of the Total Synthesis




55 %




Z:E = 7:1

Leucascandrolide A

Paterson Synthesis: Construction of C1-C8 Fragment






Paterson Synthesis: Incorporation of C11-C15 Fragment




99 %
dr 17:1

Paterson Synthesis: Further Modification of Skeleton





Paterson Synthesis: Installation of Side Chain C18-C22



Peterson Synthesis: Macrolactonization


1. TEMPO, $\text{PhI}(\text{OAc})_2$,
 CH_2Cl_2 , $20\text{ }^\circ\text{C}$
then NaClO_2 ,
 NaHPO_4 , isopentene,
aq. $t\text{BuOH}$

2. K_2CO_3 , MeOH , $20\text{ }^\circ\text{C}$




71 %

DEAD,
 PPh_3 ,
 PhH ,
 $20\text{ }^\circ\text{C}$





65 %

HF Py ,
 THF ,
 0 to $20\text{ }^\circ\text{C}$




95 %

Paterson Synthesis: Construction of Side Chain C1'-C11'




Paterson Synthesis: Completion of Total Synthesis



81 %


H₂, Lindlar,
quinoline,
EtOAc, 20 °C




92 %

Leucascandrolide A

Highlight of Other Interesting Syntheses: Kozmin




Highlight of Other Interesting Syntheses: Rychnovsky




1. Me₃OBF₄,
proton sponge,
4 Å sieves,
CH₂Cl₂, 79 %

2. OsO₄, NMO,
then NaIO₄, 80 %

3. L-Selectride,
THF, -90 to
-60 °C, 82 %



Highlight of Other Interesting Syntheses: Crimmins







Summary

- Leighton: Most linear synthesis.
- Richnovsky: Featuring Aldol-Prins cyclization to construct the skeleton.
- Wipf: Dithiane-iodide coupling was used to construct the skeleton.
- Kozmin: Most distinct feature: Spontaneous Macroacetalization.
- Carreira: Most convergent of all syntheses. Both substrate control and reagent control were used to carry out a highly diastereoselective synthesis.
- Paterson: Only substrate control of stereochemistry after the first chiral induction.
- Crimmins: Featuring selective opening of functionalized siproketal to construct the skeleton.

Rychnovsky Synthesis: Syntheses of Building Blocks



Crimmins Synthesis: Construction of Spiroketal



Crimmins Synthesis: Modification of Spiroketal

